


1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

(10) International Publication Number
WO 03/080574 A1

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(57) Abstract: The invention provides a compound of formula (I) wherein A, R¹, R², R³, R^{3a}, R⁴, R^{4a}, R⁵, and R⁶ are as defined; or a pharmaceutically acceptable salt thereof or a solvate thereof; compositions containing these compounds, processes for preparing them and their use as modulators of chemokine activity (for example CCR5 activity).



WO 03/080574 A1

Piperidine or 8-aza-bicyclo[3.2.1]oct-3-yl derivatives useful as modulators of chemokine receptor activity (especially CCR5)

The present invention relates to piperidine derivatives having pharmaceutical activity, to processes for preparing such derivatives, to pharmaceutical compositions comprising such derivatives and to the use of such derivatives as active therapeutic agents.

Pharmaceutically active piperidine derivatives are disclosed in PCT/SE01/01053, EP-A1-1013276, WO00/08013, WO01/90106, WO99/38514 and WO99/04794.

Chemokines are chemotactic cytokines that are released by a wide variety of cells to attract macrophages, T cells, eosinophils, basophils and neutrophils to sites of inflammation and also play a rôle in the maturation of cells of the immune system. Chemokines play an important rôle in immune and inflammatory responses in various diseases and disorders, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterised by a conserved four cysteine motif. The chemokine superfamily can be divided into two main groups exhibiting characteristic structural motifs, the Cys-X-Cys (C-X-C, or α) and Cys-Cys (C-C, or β) families. These are distinguished on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues and sequence similarity.

The C-X-C chemokines include several potent chemoattractants and activators of neutrophils such as interleukin-8 (IL-8) and neutrophil-activating peptide 2 (NAP-2).

The C-C chemokines include potent chemoattractants of monocytes and lymphocytes but not neutrophils such as human monocyte chemotactic proteins 1-3 (MCP-1, MCP-2 and MCP-3), RANTES (Regulated on Activation, Normal T Expressed and Secreted), eotaxin and the macrophage inflammatory proteins 1 α and 1 β (MIP-1 α and MIP-1 β).

Studies have demonstrated that the actions of the chemokines are mediated by subfamilies of G protein-coupled receptors, among which are the receptors designated CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10, CXCR1, CXCR2, CXCR3 and CXCR4. These receptors represent good targets for drug development since agents which modulate these receptors would be useful in the treatment of disorders and diseases such as those mentioned above.


The CCR5 receptor is expressed on T-lymphocytes, monocytes, macrophages, dendritic cells, microglia and other cell types. These detect and respond to several

chemokines, principally "regulated on activation normal T-cell expressed and secreted" (RANTES), macrophage inflammatory proteins (MIP) MIP-1 α and MIP-1 β and monocyte chemoattractant protein-2 (MCP-2).

This results in the recruitment of cells of the immune system to sites of disease. In many diseases it is the cells expressing CCR5 which contribute, directly or indirectly, to tissue damage. Consequently, inhibiting the recruitment of these cells is beneficial in a wide range of diseases.

CCR5 is also a co-receptor for HIV-1 and other viruses, allowing these viruses to enter cells. Blocking the receptor with a CCR5 antagonist or inducing receptor internalisation with a CCR5 agonist protects cells from viral infection.

The present invention provides a compound of formula (I):



wherein:

A is CH₂CH₂ or A is absent;

R¹ is C₃₋₇ cycloalkyl (substituted by one or two fluorine atoms and optionally further substituted by C₁₋₄ alkyl) or N-linked heterocyclyl (substituted by one or two fluorine atoms and optionally further substituted by C₁₋₄ alkyl);

R² is C₃₋₆ alkyl or C₃₋₆ cycloalkyl, or phenyl or heteroaryl either of which is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_n(C₁₋₄ alkyl), nitro, cyano or CF₃;

R^{2a}, R⁴ and R^{4a} are, independently, hydrogen or C₁₋₄ alkyl;

R³ and R^{3a} are, independently, hydrogen or C₁₋₄ alkyl or C₁₋₄ alkoxy;

R⁵ is hydrogen, C₁₋₄ alkyl (optionally substituted by halogen, hydroxy, C₁₋₄ alkoxy, C₃₋₇ cycloalkyl, SH, C₁₋₄ alkylthio, cyano or S(O)_q(C₁₋₄ alkyl)), C₃₋₄ alkenyl, C₃₋₄ alkynyl or C₃₋₇ cycloalkyl;

R⁶ is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C₁₋₂)alkyl, heteroaryl(C₁₋₂)alkyl, phenyl(C₁₋₂ alkyl)NH or heteroaryl(C₁₋₂ alkyl)NH; wherein the phenyl and heteroaryl rings of R⁶ are optionally substituted by halo, cyano, nitro, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_mC₁₋₄ alkyl, S(O)₂NR⁷R⁸, NHS(O)₂(C₁₋₄ alkyl), NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHC(O)NH₂,

C(O)NH₂, C(O)NH(C₁₋₄ alkyl), NHC(O)(C₁₋₄ alkyl), CO₂H, CO₂(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), CF₃, CHF₂, CH₂F, CH₂CF₃ or OCF₃;

R⁷ and R⁸ are, independently, hydrogen or C₁₋₄ alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C₁₋₄

5 alkyl, C(O)H or C(O)(C₁₋₄ alkyl);

m, n and q are, independently, 0, 1 or 2;

or a pharmaceutically acceptable salt thereof or a solvate thereof.

Certain compounds of the present invention can exist in different isomeric forms (such as enantiomers, diastereomers, geometric isomers or tautomers). The present invention covers
10 all such isomers and mixtures thereof in all proportions.

Suitable salts include acid addition salts (also known as adducts) such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate. An acid addition salt is, for example, a hydrochloride.

15 The compounds of the invention may exist as solvates (such as hydrates) and the present invention covers all such solvates.

Alkyl groups and moieties contain, unless otherwise specified, for example 1-6, such as 1-4, carbon atoms. Alkyl groups and moieties are straight or branched chain and are, for example, methyl, ethyl, n-propyl or iso-propyl.

20 Alkenyl includes prop-2-en-1-yl, allyl, but-3-en-1-yl, but-1-en-1-yl or 2-methylallyl. Alkynyl includes propargyl or but-3-yn-1-yl. Alkenyl and alkynyl groups and moieties are, for example, allyl or propargyl.

Cycloalkyl contains, unless otherwise specified, for example 3-7, such as 3-6, carbon atoms. Cycloalkyl is, for example, cyclopropyl, cyclobutyl or cyclopentyl.

25 When A is present the central ring of formula (I) is a 3-substituted 8-aza-bicyclo[3.2.1]oct-8-yl ring. When A is absent the central ring of formula (I) is a 4-substituted piperidin-1-yl ring.

Heterocyclyl is a non-aromatic, monocyclic ring comprising at least one nitrogen, and, optionally, one further heteroatom selected from the group comprising nitrogen, oxygen and sulphur. Heterocyclyl includes aziridinyl, azetidiny, piperidinyl, morpholinyl,
30 thiomorpholinyl, pyrrolidinyl or piperazinyl.

Heteroaryl is an aromatic 5 or 6 membered ring comprising at least one heteroatom selected from the group comprising nitrogen, oxygen and sulphur. Heteroaryl is, for example,

pyrrolyl, imidazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, thienyl, furyl, quinolinyl, isoquinolinyl, dihydroisoquinolinyl, quinazolinyl, quinoxalinyl, indolyl, isoindolyl, benzimidazolyl, benzo[b]furyl, benzo[b]thienyl, phthalazinyl, 5 benzthiazolyl or cinnolinyl.

Phenylalkyl is, for example, benzyl, 1-(phenyl)eth-1-yl or 1-(phenyl)eth-2-yl.

Heteroarylalkyl is, for example, pyridinylmethyl, pyrimidinylmethyl or 1-(pyridinyl)eth-2-yl.

The group $S(O)_2NR^7R^8$ is, for example, $S(O)_2NH_2$, $S(O)_2NH(C_{1-4} \text{ alkyl})$, $S(O)_2N(C_{1-4} \text{ alkyl})_2$, $S(O)_2(4-C(O)H\text{-piperazin-1-yl})$ or $S(O)_2(4-C(O)CH_3\text{-piperazin-1-yl})$. 10

Phenyl(C_{1-2} alkyl)NH is, for example, benzylamino. Heteroaryl(C_{1-2} alkyl)NH is, for example, pyridinyl CH_2NH , pyrimidinyl CH_2NH or pyridinyl $CH(CH_3)NH$.

In one aspect the present invention provides a compound of formula (I) wherein R^1 is C_{3-7} cycloalkyl (substituted by 1 or 2 fluorine atoms and optionally further substituted by C_{1-4} 15 alkyl).

In another aspect of the invention R^1 is C_{3-7} cycloalkyl substituted by 2 fluorine atoms.

When R^1 includes a cycloalkyl ring that ring is, for example, cyclobutyl, cyclopentyl or cyclohexyl; and further the ring is, for example, cyclohexyl.

In a further aspect of the invention R^1 is 4,4-di-fluoro-cyclohexyl, 3,3-di-fluoro-cyclopentyl or 3,3-di-fluoro-cyclobutyl. 20

In a still further aspect of the invention R^1 is, for example, 4,4-difluorocyclohex-1-yl.

In another aspect R^1 is N-linked heterocyclyl (substituted by 1 or 2 fluorine atoms and optionally further substituted by C_{1-4} alkyl). N-Linked heterocyclyl is, for example piperidin-1-yl or pyrrolidin-1-yl. R^1 is, for example, 4-fluoro-piperidin-1-yl or 3-fluoro-pyrrolidin-1-yl. 25

When R^2 is C_{3-6} alkyl it is, for example, a butyl group (such as iso-butyl) and when it is C_{3-6} cycloalkyl it is, for example, cyclopropyl or cyclohexyl.

In yet another aspect R^2 is phenyl or 6-membered heteroaryl optionally substituted in the ortho or meta position.

In a further aspect R^2 is phenyl or 6-membered heteroaryl optionally substituted (for 30 example in the 2-, 3-, or 3- and 5- positions) by halogen or CF_3 , wherein halogen is, for example, fluorine or chlorine. For example R^2 is 3-fluorophenyl, 3-chlorophenyl, 3- CF_3 -phenyl, 4-fluorophenyl or 4- CF_3 -phenyl.

In a still further aspect R^2 is optionally substituted (for example unsubstituted or substituted in the 3-, or 3- and 5- positions) phenyl (such as optionally substituted by halo (such as chloro or fluoro), cyano, methyl, ethyl, methoxy, ethoxy or CF_3); or R^2 can additionally be phenyl optionally substituted (for example unsubstituted or mono-substituted) heteroaryl (such as optionally substituted by halo (such as chloro or fluoro), cyano, methyl, ethyl, methoxy, ethoxy or CF_3).

In another aspect R^2 is optionally substituted (for example unsubstituted or substituted in the 3-, or 3- and 5- positions) phenyl (such as optionally substituted by halo (for example chloro or fluoro)). For example R^2 is phenyl, 3-fluorophenyl, 3-chlorophenyl or 3,5-difluorophenyl.

In a further aspect R^{2a} , R^3 , R^{3a} and R^4 are all hydrogen.

In still further aspect R^{4a} is hydrogen or methyl. In another aspect R^{4a} is hydrogen. In a further aspect R^{4a} is methyl.

In another aspect R^5 is hydrogen, methyl or ethyl. In yet another aspect of the invention R^5 is ethyl.

In a further aspect R^5 is iso-propyl.

In a still further aspect R^5 is C_{3-4} alkenyl, C_{3-4} alkynyl, C_{3-7} cycloalkyl or C_{3-7} cycloalkyl(C_{1-4} alkyl). For example R^5 is allyl, propargyl, cyclopropyl or cyclopropyl CH_2 . In another aspect R^5 is cyclopropyl or, for example, allyl.

In yet another aspect of the invention R^6 is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C_{1-2})alkyl, heteroaryl(C_{1-2})alkyl, phenyl(C_{1-2} alkyl)NH or heteroaryl(C_{1-2} alkyl)NH; wherein the phenyl and heteroaryl rings of R^6 are substituted by one of: $S(O)_mC_{1-4}$ alkyl, $NHC(O)NH_2$, $C(O)(C_{1-4}$ alkyl), CHF_2 , CH_2F , CH_2CF_3 or OCF_3 , and optionally further substituted by one or more of halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_mC_{1-4}$ alkyl, $S(O)_2NR^7R^8$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl), $NHC(O)(C_{1-4}$ alkyl), CO_2H , $CO_2(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ; wherein m, R^7 and R^8 are hereinbefore defined.

In a still further aspect of the invention R^6 is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C_{1-2})alkyl, heteroaryl(C_{1-2})alkyl, phenyl(C_{1-2} alkyl)NH or heteroaryl(C_{1-2} alkyl)NH (for example phenyl or phenyl CH_2); wherein the phenyl and heteroaryl rings of R^6 are substituted by $S(O)_2C_{1-4}$ alkyl, and optionally further substituted by one or more of halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_mC_{1-4}$ alkyl, $S(O)_2NR^7R^8$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl),

NHC(O)(C₁₋₄ alkyl), CO₂H, CO₂(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), CF₃, CHF₂, CH₂F, CH₂CF₃ or OCF₃; wherein m, R⁷ and R⁸ are hereinbefore defined.

In another aspect of the invention R⁶ is phenyl(C₁₋₂ alkyl) (for example benzyl); wherein the phenyl ring of R⁶ is substituted by S(O)₂C₁₋₄ alkyl, and optionally further substituted by one or more of halo, cyano, nitro, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_mC₁₋₄ alkyl, S(O)₂NR⁷R⁸, NHS(O)₂(C₁₋₄ alkyl), NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHC(O)NH₂, C(O)NH₂, C(O)NH(C₁₋₄ alkyl), NHC(O)(C₁₋₄ alkyl), CO₂H, CO₂(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), CF₃, CHF₂, CH₂F, CH₂CF₃ or OCF₃; wherein m, R⁷ and R⁸ are hereinbefore defined.

In yet another aspect of the invention R⁶ is optionally substituted benzyl, for example benzyl singly substituted (such as in the 4-position) by S(O)₂(C₁₋₄)alkyl (such as S(O)₂CH₃) or S(O)₂NR⁷R⁸ {R⁷ and R⁸ are, independently, hydrogen or C₁₋₄ alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C₁₋₄ alkyl, C(O)H or C(O)(C₁₋₄ alkyl)} (such as S(O)₂NH₂, S(O)₂NH(CH₃), S(O)₂N(CH₃)₂, S(O)₂(4-C(O)H-piperazin-1-yl) or S(O)₂(4-C(O)CH₃-piperazin-1-yl). The 5- or 6-membered ring is, for example, morpholine, thiomorpholine, piperidine, piperazine or pyrrolidine; such as piperazine.

In another aspect of the invention R⁶ is benzyl singly substituted (such as in the 4-position) by S(O)₂(C₁₋₄)alkyl (such as S(O)₂CH₃).

In a further aspect of the invention A is absent.


In another aspect of the invention A is CH₂CH₂.

In yet another aspect R⁷ and R⁸ are, independently, hydrogen or C₁₋₄ alkyl.

In a further aspect the compound of the invention is in free base form.


In a still further aspect the present invention provides a compound of formula (I) wherein A is absent or is CH₂CH₂; R¹ is C₃₋₆ cycloalkyl disubstituted with halo (such as fluoro), heterocyclyl monosubstituted by halo (such as fluoro); heterocyclyl is, for example, piperidinyl or pyrrolidinyl; R² is phenyl or monohalophenyl or dihalophenyl, where halo is, for example, fluoro, (for example R² is phenyl, 3-fluorophenyl or 3,5-difluorophenyl); R^{2a}, R³, R^{3a} and R⁴ are all hydrogen; R^{4a} is hydrogen or C₁₋₄ alkyl (such as methyl); R⁵ is C₁₋₄ alkyl (such as ethyl); R⁶ is benzyl singly substituted (such as in the 4-position) by S(O)₂(C₁₋₄)alkyl (such as S(O)₂CH₃); or an acid addition salt thereof (such as a hydrochloride).

In yet another aspect the present invention provides a compound of formula (Ia):




wherein R¹ and R² are as defined above, and having the absolute configuration shown.

In a further aspect the present invention provides a compound of formula (Ib):



5 wherein R¹ and R² are as defined above, and having the absolute configuration shown.

In a still further aspect the present invention provides a compound of formula (Ic):




wherein R¹ and R² are as defined above, and having the absolute configuration shown.

The compounds in Tables I, II and III illustrate the invention.

TABLE I

Table I comprises compounds of the invention having the formula (Ia).




Compound No.	R ¹	R ²	Adduct	LCMS (MH ⁺)
1	4,4-difluoro-cyclohexyl	Phenyl		604
2	4-fluoro-piperidin-1-yl	Phenyl		587
3	(R)-3-fluoro-pyrrolidin-1-yl	Phenyl		573
4	(S)-3-fluoro-pyrrolidin-1-yl	Phenyl		573
5	4,4-difluoro-cyclohexyl	3-fluoro-phenyl		594
6	3,3-difluoro-cyclobutyl	3,5-difluoro-phenyl	hydrochloride	612
7	4,4-difluoro-cyclohexyl	3,5-difluoro-phenyl	hydrochloride	640
8	3,3-difluoro-cyclobutyl	Phenyl	hydrochloride	576
9	3,3-difluoro-cyclobutyl	3-fluoro-phenyl	hydrochloride	594
10	(R)-3,3-difluoro-cyclopentyl	Phenyl		
11	(S)-3,3-difluoro-cyclopentyl	Phenyl		
12	(R)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl		

13	(S)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl	
14	(R)-3,3-difluoro-cyclopentyl	3-fluoro-phenyl	
15	(S)-3,3,3-difluoro-cyclopentyl	3-fluoro-phenyl	

TABLE II

Table I comprises compounds of the invention having the formula (Ib).




Compound No.	R ¹	R ²	LCMS (MH ⁺)
1	4,4-difluoro-cyclohexyl	Phenyl	604
2	4-fluoro-piperidin-1-yl	Phenyl	
3	(R)-3-fluoro-pyrrolidin-1-yl	Phenyl	
4	(S)-3-fluoro-pyrrolidin-1-yl	Phenyl	
5	4,4-difluoro-cyclohexyl	3-fluoro-phenyl	
6	3,3-difluoro-cyclobutyl	3,5-difluoro-phenyl	
7	4,4-difluoro-cyclohexyl	3,5-difluoro-phenyl	
8	3,3-difluoro-cyclobutyl	Phenyl	
9	3,3-difluoro-cyclobutyl	3-fluoro-phenyl	

10	(R)-3,3-difluoro-cyclopentyl	Phenyl	
11	(S)-3,3-difluoro-cyclopentyl	Phenyl	
12	(R)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl	
13	(S)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl	
14	(R)-3,3-difluoro-cyclopentyl	3-fluoro-phenyl	
15	(S)-3,3-difluoro-cyclopentyl	3-fluoro-phenyl	

TABLE III

Table I comprises compounds of the invention having the formula (Ic).




Compound No.	R ¹	R ²	Adduct	LCMS (MH ⁺)
1	4,4-difluoro-cyclohexyl	Phenyl	hydrochloride	618
2	4-fluoro-piperidin-1-yl	Phenyl		
3	(R)-3-fluoro-pyrrolidin-1-yl	Phenyl		
4	(S)-3-fluoro-pyrrolidin-1-yl	Phenyl		
5	4,4-difluoro-cyclohexyl	3-fluoro-phenyl		
6	3,3-difluoro-cyclobutyl	3,5-difluoro-phenyl		

7	4,4-difluoro-cyclohexyl	3,5-difluoro-phenyl		
8	3,3-difluoro-cyclobutyl	Phenyl	hydrochloride	590
9	3,3-difluoro-cyclobutyl	3-fluoro-phenyl		
10	(<i>R</i>)-3,3-difluoro-cyclopentyl	Phenyl		
11	(<i>S</i>)-3,3-difluoro-cyclopentyl	Phenyl		
12	(<i>R</i>)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl		
13	(<i>S</i>)-3,3-difluoro-cyclopentyl	3,5-difluoro-phenyl		
14	(<i>R</i>)-3,3-difluoro-cyclopentyl	3-fluoro-phenyl		
15	(<i>S</i>)-3,3-difluoro-cyclopentyl	3-fluoro-phenyl		


The compounds of formulae (I), (Ia), (Ib) and (Ic) can be prepared as described below, by adaptation of methods described in the art (such as WO 01/90106) or by following or adapting the Examples or Methods provided below.

Specifically, a compound of formula (I) or (Ia) can be prepared by treating a
5 compound of formula (II):




with: an acid chloride of formula $R^1C(O)Cl$, in the presence of a base (such as a tertiary amine, for example triethylamine) and in a suitable solvent (such as a chlorinated hydrocarbon, for example dichloromethane); or an acid of formula R^1CO_2H in the presence of
10 a suitable coupling agent (such as O-(7-azabenzotriazol-1-yl)- N,N,N',N' -tetramethyluronium hexafluorophosphate [HATU] or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate [PyBrop]) in the presence of a suitable base (such as a tertiary amine, for example diisopropylethylamine) in a suitable solvent (such as N -methylpyrrolidinone).

A compound of formula (II) can be prepared by treating a compound of formula (III):




15 with trifluoroacetic acid or hydrochloric acid in the presence of methanol, and then basifying to release the free amine form of formula (II).

A compound of formula (III) can be prepared by reductively aminating a compound of formula (IV):




20 with a compound of formula (V):



in the presence of a suitable solvent (such as an aliphatic alcohol such as methanol), a suitable organic acid (such as an aliphatic acid, for example acetic acid) and a suitable reducing agent (such as sodium triacetoxyborohydride or sodium cyanoborohydride).

- 5 A compound of formula (II) wherein R^{2a} is hydrogen can be prepared by reductive amination of a compound of formula (VI):



- for example by reacting a compound of formula (VI) with hydroxylamine and hydrogenating the product so formed with hydrogen in the presence of a suitable metal catalyst (such as palladium or platinum catalyst, for example palladium on charcoal).
- 10

A compound of formula (VI), wherein R^{4a} is hydrogen, can be prepared by reacting a compound of formula (V) with:

- an alkyl halide of formula $R^2C(O)CR^3R^{3a}CHR^4X$ (wherein X is halogen, such as chloro, bromo or iodo) in the presence of a suitable base (such as potassium carbonate) and a suitable solvent (such as acetone); or,
 - compounds of formula $R^2C(O)CHR^3R^{3a}$ and R^4CHO in the presence of a suitable acid (such as acetic acid).
- 15

A compound of formula (VI), wherein R^{3a} is hydrogen, can be prepared by reacting a compound of formula (V) with an alkene of formula $R^2C(O)CR^3=CR^4R^{4a}$ in a suitable solvent (such as an aliphatic alcohol, for example ethanol) at a temperature in the range -10 to 100°C.

20

Compounds of formula (Ib) can be prepared by referring to WO 01/90106 and WO 01/87839.

- The starting materials for these processes are commercially available, can be prepared by literature methods or can be prepared by adapting literature methods. In a further aspect the invention provides processes for preparing the compounds of formulae (I), (Ia), (Ib) and
- 25

(Ic). Many of the intermediates in the processes are novel and these are provided as further features of the invention.

The compounds of the invention have activity as pharmaceuticals, in particular as modulators (such as agonists, partial agonists, inverse agonists or antagonists) of chemokine receptor (for example CCR5) activity, and may be used in the treatment of autoimmune, inflammatory, proliferative or hyperproliferative diseases, or immunologically-mediated diseases (including rejection of transplanted organs or tissues and Acquired Immunodeficiency Syndrome (AIDS)). Examples of these conditions are:

- (1) (the respiratory tract) obstructive diseases of airways including: chronic obstructive pulmonary disease (COPD) (such as irreversible COPD); pulmonary fibrosis; asthma {such as bronchial, allergic, intrinsic, extrinsic or dust asthma, particularly chronic or inveterate asthma (for example late asthma or airways hyper-responsiveness)}; bronchitis {such as eosinophilic bronchitis}; acute, allergic, atrophic rhinitis or chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca or rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous or pseudomembranous rhinitis or scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) or vasomotor rhinitis; sarcoidosis; farmer's lung and related diseases; nasal polyposis; fibroid lung or idiopathic interstitial pneumonia;
- (2) (bone and joints) arthrides including rheumatic, infectious, autoimmune, seronegative spondyloarthropathies (such as ankylosing spondylitis, psoriatic arthritis or Reiter's disease), Behcet's disease, Sjogren's syndrome or systemic sclerosis;
- (3) (skin and eyes) psoriasis, atopic dermatitis, contact dermatitis or other eczematous dermatides, seborrhoetic dermatitis, lichen planus, pemphigus, bullous pemphigus, epidermolysis bullosa, urticaria, angiodermas, vasculitides erythemas, cutaneous eosinophilias, uveitis, alopecia areata or vernal conjunctivitis;
- (4) (gastrointestinal tract) Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, irritable bowel disease or food-related allergies which have effects remote from the gut (for example migraine, rhinitis or eczema);
- (5) (Allograft rejection) acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin or cornea; or chronic graft versus host disease; and/or
- (6) (other tissues or diseases) Alzheimer's disease, multiple sclerosis, atherosclerosis, inhibiting the entry of viruses into target cells, Acquired Immunodeficiency Syndrome

(AIDS), lupus disorders (such as lupus erythematosus or systemic lupus), erythematosus, Hashimoto's thyroiditis, myasthenia gravis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, leprosy (such as lepromatous leprosy), Peridontal disease, sezary syndrome, idiopathic thrombocytopenia pupura, disorders of the menstrual cycle,
5 glomerulonephritis or cerebral malaria.

The compounds of the present invention are also of value in inhibiting the entry of viruses (such as human immunodeficiency virus (HIV)) into target cells and, therefore, are of value in the prevention of infection by viruses (such as HIV), the treatment of infection by viruses (such as HIV) and the prevention and/or treatment of acquired immune deficiency
10 syndrome (AIDS).

According to a further feature of the invention there is provided a compound of the formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof or a solvate thereof, for use in a method of treatment of a warm blooded animal (such as man) by therapy (including prophylaxis).

15 According to a further feature of the present invention there is provided a method for modulating chemokine receptor activity (for example CCR5 receptor activity) in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof or a solvate thereof.

20 The present invention further provides a method of treating a chemokine mediated disease state (for example a CCR5 mediated disease state, such as rheumatoid arthritis) in a warm blooded animal (such as man) suffering from, or at risk of, said disease, which comprises administering to an animal in need of such treatment a therapeutically effective amount of a compound of formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula
25 (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof or solvate thereof.

The invention also provides a compound of the formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof or a solvate thereof, for use in therapy (including prophylaxis); for example in the treatment of a chemokine mediated disease state (for example a CCR5 mediated disease state) in a warm
30 blooded animal, such as man, such as in the treatment of rheumatoid arthritis.

The invention also provides a compound of the formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof

or a solvate thereof, for use as a medicament, for example a medicament for the treatment of rheumatoid arthritis.

In another aspect the present invention provides the use of a compound of the formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof or a solvate thereof, in the manufacture of a medicament for use in therapy (for example in modulating chemokine receptor activity (for example CCR5 receptor activity (for example in the treatment of rheumatoid arthritis)) in a warm blooded animal, such as man).

The invention further provides the use of a compound of formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of:

- (1) (the respiratory tract) obstructive diseases of airways including: chronic obstructive pulmonary disease (COPD) (such as irreversible COPD); asthma {such as bronchial, allergic, intrinsic, extrinsic or dust asthma, particularly chronic or inveterate asthma (for example late asthma or airways hyper-responsiveness)}; bronchitis {such as eosinophilic bronchitis}; acute, allergic, atrophic rhinitis or chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca or rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous or pseudomembranous rhinitis or scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) or vasomotor rhinitis; sarcoidosis; farmer's lung and related diseases; nasal polyposis; fibroid lung or idiopathic interstitial pneumonia;
- (2) (bone and joints) arthrides including rheumatic, infectious, autoimmune, seronegative spondyloarthropathies (such as ankylosing spondylitis, psoriatic arthritis or Reiter's disease), Behcet's disease, Sjogren's syndrome or systemic sclerosis;
- (3) (skin and eyes) psoriasis, atopic dermatitis, contact dermatitis or other eczematous dermatides, seborrhoetic dermatitis, lichen planus, pemphigus, bullous pemphigus, epidermolysis bullosa, urticaria, angiodermas, vasculitides erythemas, cutaneous eosinophilias, uveitis, alopecia areata or vernal conjunctivitis;
- (4) (gastrointestinal tract) Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, irritable bowel disease or food-related allergies which have effects remote from the gut (for example migraine, rhinitis or eczema);
- (5) (Allograft rejection) acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin or cornea; or chronic graft versus host disease; and/or

(6) (other tissues or diseases) Alzheimer's disease, multiple sclerosis, atherosclerosis, Acquired Immunodeficiency Syndrome (AIDS), lupus disorders (such as lupus erythematosus or systemic lupus), erythematosus, Hashimoto's thyroiditis, myasthenia gravis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, leprosy (such as lepromatous leprosy), Peridontal disease, sezary syndrome, idiopathic thrombocytopenia pupura or disorders of the menstrual cycle;
in a warm blooded animal, such as man.

In order to use a compound of the invention, or a pharmaceutically acceptable salt thereof or solvate thereof, for the therapeutic treatment of a warm blooded animal, such as man, in particular modulating chemokine receptor (for example CCR5 receptor) activity, said ingredient is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

Therefore in another aspect the present invention provides a pharmaceutical composition which comprises a compound of the formula (I), (Ia), (Ib) or (Ic) (for example a compound of formula (I), (Ia) or (Ib)), or a pharmaceutically acceptable salt thereof or a solvate thereof (active ingredient), and a pharmaceutically acceptable adjuvant, diluent or carrier. In a further aspect the present invention provides a process for the preparation of said composition which comprises mixing active ingredient with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will, for example, comprise from 0.05 to 99 %w (per cent by weight), such as from 0.05 to 80 %w, for example from 0.10 to 70 %w, such as from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

The pharmaceutical compositions of this invention may be administered in standard manner for the disease condition that it is desired to treat. A suitable pharmaceutical composition of this invention is one suitable for oral administration in unit dosage form, for example a tablet or capsule which contains between 0.1mg and 1g of active ingredient.

Each patient may receive, for example, an intravenous, subcutaneous or intramuscular dose of 0.01mgkg^{-1} to 100mgkg^{-1} of the compound, for example in the range of 0.1mgkg^{-1} to 20mgkg^{-1} of this invention, the composition being administered 1 to 4 times per day. The intravenous, subcutaneous and intramuscular dose may be given by means of a bolus injection. Alternatively the intravenous dose may be given by continuous infusion over a period of time. Alternatively each patient will receive a daily oral dose which is

approximately equivalent to the daily parenteral dose, the composition being administered 1 to 4 times per day.

The invention will now be illustrated by the following non-limiting Examples in which, unless stated otherwise:

- 5 (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mm Hg) with a bath temperature of up to 60°C;
- 10 (iii) chromatography unless otherwise stated means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column is referred to, this means a column containing 10g or 20g of silica of 40 micron particle size, the silica being contained in a 60ml disposable syringe and supported by a porous disc, obtained from Varian, Harbor City, California, USA under the name "Mega Bond Elut SI".
- 15 Where an "IsoluteTM SCX column" is referred to, this means a column containing benzenesulphonic acid (non-endcapped) obtained from International Sorbent Technology Ltd., 1st House, Duffryn Industrial Estate, Ystrad Mynach, Hengoed, Mid Glamorgan, UK. Where "ArgonautTM PS-*tris*-amine scavenger resin" is referred to, this means a *tris*-(2-aminoethyl)amine polystyrene resin obtained from Argonaut Technologies Inc., 887 Industrial
- 20 Road, Suite G, San Carlos, California, USA.
- (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) yields, when given, are for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was
- 25 required;
- (vi) when given, ¹H NMR data is quoted and is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio DMSO (CD₃SOCD₃) as the solvent unless otherwise stated; coupling constants (J) are given in Hz;
- 30 (vii) chemical symbols have their usual meanings; SI units and symbols are used;
- (viii) solvent ratios are given in percentage by volume;

(ix) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionisation (APCI) mode using a direct exposure probe; where indicated ionisation was effected by electrospray (ES); where values for m/z are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the positive mass ion - $(M+H)^+$;

(x) LCMS characterisation was performed using a pair of Gilson 306 pumps with Gilson 233 XL sampler and Waters ZMD4000 mass spectrometer. The LC comprised water symmetry 4.6x50 column C18 with 5 micron particle size. The eluents were: A, water with 0.05% formic acid and B, acetonitrile with 0.05% formic acid. The eluent gradient went from 95% A to 95% B in 6 minutes. Where indicated ionisation was effected by electrospray (ES); where values for m/z are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the positive mass ion - $(M+H)^+$ and

(xi) the following abbreviations are used:

THF	tetrahydrofuran;
Boc	tert-butoxycarbonyl;
THF	tetrahydrofuran;
DCM	dichloromethane; and
HATU	O-(7-Azabenzotriazol-1-yl)- N,N,N',N' -tetramethyluronium hexafluorophosphate.

EXAMPLE 1

This Example illustrates the preparation of (*S*)-*N*-[1-(3-phenyl-3-[4,4-difluorocyclohexylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (Compound No. 1 of Table I).

(*S*)-*N*-[1-(3-Phenyl-3-aminopropyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide dihydrochloride (Method A, 250mg), 4,4-difluorocyclohexane carboxylic acid (100mg) and *N,N*-di-isopropylethylamine (0.7mL) were stirred in DCM (5mL) at room temperature. To this solution was added HATU (200mg) and stirring was continued for 16 hours. 2N Sodium hydroxide solution (2mL) was added and the organic layer separated, washed with water and concentrated; the residue was purified by silica gel chromatography (eluent 0-30% methanol in ethyl acetate) to give the title compound as a colourless gum (110mg); NMR: 1.0 and 1.1 (t, 3H), 1.7 (m, 7H), 2.2 (m, 6H), 3.0 (m,

3H), 3.2 (s, 3H), 3.4 (q, 2H), 3.8 and 3.9 (s, 2H), 4.1 and 4.3 (m, 1H), 4.8 (m, 1H), 7.2 (m, 1H), 7.3 (m, 4H), 7.5 (d, 2H), 7.8 (d, 2H), 8.85 (m, 1H); MS: 604 (MH⁺).

The procedure described in Example 1 can be repeated using different carboxylic acids (such as 3,3-di-fluorocyclobutane carboxylic acid) in place of 4,4-difluorocyclohexane carboxylic acid and different amines or amine dihydrochlorides (such as (*S*)-*N*-{1-[3-amino-3-(3-fluorophenyl)propyl]piperidin-4-yl}-*N*-ethyl-2-(4-methanesulfonyl-phenyl)acetamide (Method F), (*S*)-*N*-{1-[3-amino-3-(3,5-di-fluorophenyl)propyl]piperidin-4-yl}-*N*-ethyl-2-(4-methanesulfonyl-phenyl)acetamide dihydrochloride (Method G)) or *N*-[1-((4*S*)-4-phenyl-4-aminobut-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide dihydrochloride (Method H)) in place of (*S*)-*N*-[1-(3-phenyl-3-aminopropyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide dihydrochloride.

EXAMPLE 2

This Example illustrates the preparation of (*S*)-*N*-[1-(3-phenyl-3-[4-fluoropiperidin-1-ylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (Compound No. 2 of Table I).

To (*S*)-*N*-[1-(3-phenyl-3-[4-nitrophenoxy-carboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (Method C, 150mg) in DCM (10mL) was added 4-fluoropiperidine hydrochloride (100mg) and *N,N*-di-isopropylethylamine (1mL). The resulting mixture was stirred at room temperature for 16 hours. 2N Sodium hydroxide solution (10mL) was added and the organic layer separated, washed with water, dried (MgSO₄) and concentrated; the residue was purified by silica gel chromatography (eluent 0-20% methanol in ethyl acetate) to give the title compound as a colourless gum (140mg); MS: 587 (MH⁺).

The procedure described in Example 2 can be repeated using different amines (such as (*S*)-3-fluoro-pyrrolidine or (*R*)-3-fluoropyrrolidine) in place of 4-fluoropiperidine hydrochloride.

EXAMPLE 3

This Example illustrates the preparation of (*S*)-4,4-difluoro-cyclohexanecarboxylic acid [3-(3-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-8-aza-bicyclo[3.2.1]oct-8-yl-exo)-1-phenyl-propyl]-amide (Compound No. 1 of Table II).

To a solution of *N*-(8-aza-bicyclo[3.2.1]oct-3-yl-exo)-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide (Method D; 98mg, 0.28mmol) in DCM was added (*S*)-3-phenyl-3-(4,4-difluorocyclohexylcarbonylamino)propanal (Method E; 165mg, 0.56mmol). To the resulting mixture was added sodium triacetoxyborohydride (119mg). This was then stirred at room temperature for 18 h, washed with water, dried over MgSO₄ and concentrated. Purification was achieved by BondElut chromatography eluting with a gradient of DCM to 10% methanol and 1% 0.88 ammonia in DCM to give the title compound (143mg); NMR (CDCl₃): 1.1 and 1.2 (t, 3H), 1.3 (m, 1H), 1.9 (m, 19H), 2.3 (m, 1H), 2.5 (m, 1H), 3.0 (s, 3H), 3.3 (m, 4H), 3.8 (m, 2H), 3.6 and 4.4 (m, 1H), 5.0 (m, 1H), 7.2 (m, 5H), 7.4 (m, 2H), 7.9 (m, 2H); MS: 630 (MH⁺).

Below are presented certain NMR data for some compounds of the invention.

(*S*)-*N*-[1-(3-Phenyl-3-[(*R*)-3-fluoropyrrolidin-1-ylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (Compound No. 3 of Table I).

NMR (d₆-DMSO, 120°C): 1.1 (t, 3H), 1.5 (m, 2H), 1.8 (m, 2H), 1.9 (m, 2H), 2.1 (m, 2H), 2.3 (m, 2H), 2.9 (m, 2H), 3.15 (s, 3H), 3.3 (m, 2H), 3.35 (m, 2H), 3.5 (m, 2H), 3.6 (dd, 1H), 3.8 (s, 2H), 3.85 (m, 1H), 4.9 (dd, 1H), 5.3 (d, 1H), 6.25 (d, 1H), 7.2 (m, 1H), 7.3 (m, 4H), 7.55 (d, 2H), 7.85 (d, 2H).

(*S*)-*N*-[1-(3-Phenyl-3-[(*S*)-3-fluoropyrrolidin-1-ylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (Compound No. 4 of Table I).

NMR (d₆-DMSO, 120°C): 1.1 (t, 3H), 1.5 (m, 2H), 1.8 (m, 2H), 1.9 (m, 2H), 2.1 (m, 2H), 2.3 (m, 2H), 2.9 (m, 2H), 3.15 (s, 3H), 3.3 (m, 2H), 3.35 (m, 2H), 3.5 (m, 2H), 3.6 (dd, 1H), 3.8 (s, 2H), 3.85 (m, 1H), 4.9 (dd, 1H), 5.3 (d, 1H), 6.25 (d, 1H), 7.2 (m, 1H), 7.3 (m, 4H), 7.55 (d, 2H), 7.85 (d, 2H).

(*S*)-*N*-[1-(3-[3,5-Difluorophenyl]-3-[3,3-difluorocyclobutylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 6 of Table I).

NMR (d6-DMSO, 120°C): 1.15 (t, 3H), 1.8 (m, 2H), 2.3 (m, 2H), 2.4 (m, 2H), 2.7-3.0 (m, 9H), 3.15 (s, 3H), 3.35 (q, 2H), 3.45 (m, 2H), 3.85 (s, 2H), 4.2 (br m, 1H), 5.0 (dd, 1H), 6.95 (dd, 1H), 7.1 (d, 2H), 7.5 (d, 2H), 7.85 (d, 2H), 8.45 (br s, 1H).

5 *(S)-N*-[1-(3-[3,5-Difluorophenyl]-3-[4,4-difluorocyclohexylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 7 of Table I).

10 NMR (d6-DMSO, 120°C): 1.1 (t, 3H), 1.7 (m, 8H), 2.1 (m, 2H), 2.3 (m, 2H), 2.4 (m, 4H), 3.0 (m, 3H), 3.15 (s, 3H), 3.35 (q, 2H), 3.45 (m, 2H), 3.85 (s, 2H), 4.2 (br m, 1H), 4.9 (m, 1H), 6.9 (dd, 1H), 7.05 (d, 2H), 7.5 (d, 2H), 7.85 (d, 2H), 8.3 (br s, 1H).

(S)-N-[1-(3-Phenyl-3-[3,3-difluorocyclobutylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 8 of Table I).

15 NMR (d6-DMSO, 120°C): 1.2 (t, 3H), 1.8 (m, 2H), 2.3 (m, 2H), 2.4 (m, 2H), 2.75 (m, 4H), 3.05 (m, 5H), 3.2 (s, 3H), 3.4 (q, 2H), 3.45 (m, 2H), 3.9 (s, 2H), 4.2 (br m, 1H), 5.0 (dd, 1H), 7.3 (m, 1H), 7.35 (m, 4H), 7.55 (d, 2H), 7.95 (d, 2H), 8.35 (br s, 1H).

20 *(S)-N*-[1-(3-[3-Fluorophenyl]-3-[3,3-difluorocyclobutylcarboxyamino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 9 of Table I).

 NMR (d6-DMSO, 120°C): 1.15 (t, 3H), 1.8 (m, 2H), 2.25 (m, 2H), 2.35 (m, 2H), 2.7-3.0 (m, 9H), 3.2 (s, 3H), 3.4 (q, 2H), 3.45 (m, 2H), 3.9 (s, 2H), 4.2 (br m, 1H), 5.0 (dd, 1H), 7.05 (m, 1H), 7.25 (m, 2H), 7.35 (m, 1H), 7.55 (d, 2H), 7.9 (d, 2H), 8.35 (br s, 1H).

25 *N*-[1-((4*S*)-4-Phenyl-4-[4,4-difluorocyclohexylcarboxyamino]but-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 1 of Table III).

30 NMR: 1.1 & 1.23 (t, 3H), 1.46 (d, 3H), 1.60 (m, 2H), 1.67 (m, 2H), 1.80 (m, 2H), 1.97 (m, 1H), 2.34 - 2.60 (m, 4H), 3.14 (m, 3H), 3.24 (s, 3H), 3.24 - 3.49 (m, 6H), 3.90 (m, 4H), 4.26 (m, 1H), 5.03 (m, 1H), 7.32 (m, 1H), 7.38 (m, 2H), 7.48 (m, 2H), 7.57 (m, 2H), 7.91 (dd, 2H), 8.48 (t, 1H).


N-[1-((4*S*)-4-Phenyl-4-[3,3-difluorocyclobutylcarboxyamino]but-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide hydrochloride (Compound No. 8 of Table III).

NMR (d6-DMSO, 120°C): 1.15 (t, 3H), 1.3 (d, 3H), 1.8 (m, 2H), 2.15 (m, 1H), 2.55 (m, 2H), 2.75 (m, 5H), 3.1-3.2 (m, 3H), 3.2 (s, 3H), 3.3 (m, 2H), 3.4 (q, 2H), 3.55 (m, 1H), 3.9 (s, 2H), 4.3 (br m, 1H), 5.0 (dd, 1H), 7.3 (m, 1H), 7.35 (m, 2H), 7.45 (m, 2H), 7.55 (d, 2H), 7.9 (d, 2H), 8.3 (br s, 1H).


5

Method A

(S)-N-[1-(3-Phenyl-3-aminopropyl)-4-piperidinyl]-N-ethyl-4-methanesulfonylphenylacetamide dihydrochloride




10 Step 1: Preparation of 1-phenylmethyl-4-ethylaminopiperidine dihydrochloride



To a solution of 1-phenylmethyl-4-piperidone (25.0 g, 132 mmol) in THF (250 mL) was added ethylamine hydrochloride (12.0 g, 147 mmol) and methanol (50 mL) and the resulting mixture stirred at room temperature for 10 min. Sodium triacetoxyborohydride (40 g, 189 mmol) was added portionwise and the resulting mixture stirred at room temperature for 1 h. 2M Sodium hydroxide solution (250 mL) was added and the resulting mixture extracted with diethyl ether. The organic extracts were dried (K₂CO₃) and evaporated to give 1-phenylmethyl-4-ethylaminopiperidine as an oil. This was dissolved in ethanol (500 mL) and concentrated hydrochloric acid (20 mL) was added. The resulting crystals were collected, washed with diethyl ether and dried giving the sub-titled compound as a solid (38 g); NMR: (CDCl₃): 1.10 (t, 3H), 1.40 (m, 2H), 1.83 (m, 2H), 2.02 (m, 2H), 2.65 (q, 2H), 2.85 (m, 2H), 3.50 (s, 2H), 3.75 (m, 1H), 7.2 - 7.4 (m, 5H); MS: 219 (MH⁺).


20

Step 2: Preparation of *N*-(1-phenylmethyl-4-piperidiny)-*N*-ethyl-4-methanesulfonylphenylacetamide



To a solution of 1-phenylmethyl-4-ethylaminopiperidine dihydrochloride (32.0g, 110mmol) in DCM (500mL) was added *N,N*-diisopropylethylamine (60mL) with stirring to ensure complete dissolution. 4-Methanesulfonylphenylacetic acid (25.0g, 117mmol), 4-Dimethylaminopyridine (4-DMAP) (2.0g) and dicyclohexylcarbodiimide (DCCI) (25.0g, 121mmol) were added and the resulting mixture was stirred at room temperature for 20 h. The precipitate was removed by filtration and the resulting solution was washed successively with 2N aqueous HCl, water and 1N aqueous NaOH, dried (MgSO₄) and evaporated. The residue was purified by silica gel chromatography (eluent 10% MeOH/ethyl acetate) to afford the sub-titled compound (35 g, 76%); NMR: 1.00 and 1.14 (t, 3H), 1.45 and 1.70 (m, 2H), 1.95 (br m, 2H), 2.80 (br m, 2H), 3.18 (s, 3H), 3.20 and 3.33 (q, 2H), 3.45 (s, 2H), 3.80 and 3.87 (s, 2H), 3.70 and 4.10 (m, 1H), 7.2 - 7.3 (m, 5H), 7.48 (m, 2H), 7.82 (m, 2H); MS: 415 (MH⁺).

Step 3: Preparation of *N*-(4-Piperidiny)-*N*-ethyl-4-methanesulfonylphenylacetamide




To a solution of *N*-(1-phenylmethyl-4-piperidiny)-*N*-ethyl-4-methanesulfonylphenylacetamide (34g, 82mmol) in ethanol (600mL) was added ammonium formate (40g). The mixture was purged with argon and 30% Pd on carbon (4.2g) was added. The resulting mixture was stirred at reflux for 4 h, then allowed to cool and filtered through diatomaceous earth. The filtrate was evaporated to give a thick oil which solidified on standing to yield the sub-titled compound (24.9 g, 94%); NMR: 1.02 and 1.15 (t, 3H), 1.4 - 1.6 (br m, 4H), 2.45 (m, 2H), 2.93 (br m, 2H), 3.18 (s, 3H), 3.20 and 3.32 (q, 2H), 3.72 and 4.18 (m, 1H), 3.80 and 3.87 (s, 2H), 7.50 (m, 2H), 7.85 (m, 2H); MS: 325 (MH⁺).

Step 4: Preparation of title compound


To a solution of (*S*)-3-phenyl-3-Bocaminopropanal (Method B, 1.4g, 5.6mmol) in ethanol (100mL) and DCM (50mL) was added *N*-(4-piperidiny)-*N*-ethyl-4-methanesulfonylphenylacetamide (2.0g, 6.2mmol), glacial acetic acid (0.6mL, 10mmol) and sodium triacetoxyborohydride (2.0g, 9.4mmol) and the resulting mixture was stirred at room temperature for 18 h. The mixture was partitioned between DCM and 2M aqueous sodium hydroxide (35mL), and the organic phase was washed with water, dried and concentrated. The residue was suspended in methanol (10mL) and concentrated hydrochloric acid (10mL) was added. The resulting mixture was stirred for 30 minutes then evaporated. The residue was azeotroped with ethanol and toluene and triturated with diethyl ether yielding the title compound as a solid (1.3g); NMR (d6 DMSO at 373K): 1.1 (t, 3H), 1.5 (m, 2H), 1.9 (m, 2H), 2.0 (m, 1H), 2.3 (m, 2H), 3.0 (m, 1H), 3.2 (m, 4H), 3.3 (q, 2H), 3.9 (s, 2H), 4.0 (m, 1H), 4.4 (m, 1H), 7.4 (m, 3H), 7.5 (m, 4H), 7.9 (m, 2H); MS: 458.

15 Method B

(*S*)-3-Phenyl-3-Boc-aminopropanal



Step 1: Preparation of (*S*)-*N*-Methyl-*N*-methoxy-3-phenyl-3-Bocaminopropionamide



20 To a solution of (*S*)-3-phenyl-3-Bocaminopropanoic acid (available from PepTech Corp. of Cambridge, Massachusetts, USA; 4.97g, 18.7mmol) in DCM (100mL) was added DIPEA (14.8mL, 84.8mmol) and *N,O*-dimethylhydroxylamine hydrochloride (2.21g, 22.7mmol) followed by HATU (8.44g, 84.8mmol). The resulting mixture was stirred at room temperature for 18h, diluted with DCM, washed with 2M aqueous sodium hydroxide and water. The organic phase was dried (Na₂SO₄) and concentrated. The residue was purified by silica column chromatography (eluting with isohexane then 3:1 ethyl acetate to isohexane)

25


giving the sub-titled compound as a colourless oil (5.58g, 97%); NMR (CDCl₃): 1.40 (s, 9H), 2.83 (dd, 1H), 3.01 (m, 1H), 3.08 (s, 3H), 3.52 (s, 3H), 5.10 (m, 1H), 7.28 (m, 5H); MS: 309.

Step 2: Preparation of title compound

5 To a solution of (*S*)-*N*-methyl-*N*-methoxy-3-phenyl-3-Bocaminopropionamide (17.9mmol) in toluene (180mL) at -20°C was added sodium bis(2-methoxyethoxy)aluminium hydride (65% solution in toluene, 35.8mmol) dropwise. The resulting mixture was stirred at -15°C for 1h. The mixture was washed with saturated aqueous sodium dihydrogen phosphate solution (250mL). The organic phase was dried (Na₂SO₄) and concentrated to give the title
10 compound (5g); NMR: 1.4 (s, 9H), 2.8 (m, 2H), 5.1 (m, 1H), 7.3 (m, 5H), 8.6 (m, 1H), 9.6 (t, 1H).

Method C


(*S*)-*N*-[1-(3-phenyl-3-[4-nitrophenoxy-carboxy-amino]propyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide
15



To (*S*)-*N*-[1-(3-phenyl-3-aminopropyl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide dihydrochloride (2.0g, 3.8mmol) in DCM (50mL) was added *N,N*-di-isopropylethylamine (2mL) and 4-nitrophenyl chloroformate (1.0g, 4.9mmol)
20 and the resulting mixture stirred at ambient temperature for 16 hours. The mixture was washed with saturated sodium bicarbonate solution (50mL) and dried over anhydrous magnesium sulphate. The residue was purified by silica gel chromatography (eluent 0-10% methanol in ethyl acetate) to give the title compound as a pale yellow gum (2g).


Method D

N-(8-Aza-bicyclo[3.2.1]oct-3-yl-exo)-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide



Step 1: Preparation of *N*-(8-Benzyl-8-aza-bicyclo[3.2.1]oct-3-yl)-*N*-ethyl-2-(4-

5 methanesulfonyl-phenyl)-acetamide



To a solution of 8-Benzyl-8-aza-bicyclo[3.2.1]oct-3-yl-exo-amine (John S. Kiely, Marland P. Hutt, Townley P. Culbertson, Ruth A. Bucsh and Donald F. Worth; J. Med. Chem., 1991, **34**, 656; 2.81g, 13 mmol) in DCM (40mL) was added acetaldehyde (0.69g, 16mmol) and the resulting mixture stirred at room temperature for 1h. Sodium triacetoxyborohydride (3.3 g, 16mmol) was added portionwise and the resulting mixture stirred at room temperature for 16h. The mixture was then washed with water, dried over MgSO₄ and concentrated. This material was then dissolved in DCM (50mL) and 4-methanesulfonylphenylacetic acid (3.1g, 14mmol) and diisopropylcarbodiimide (2.1g, 14mmol) were added and the resulting mixture stirred for 2h. The precipitate was removed by filtration and the crude material was adsorbed onto silica. Silica gel chromatography (eluent: 100% DCM to 10% methanol and 1% 0.88 ammonia in DCM) gave the sub-titled compound as a foam (0.37g); NMR (CDCl₃): 1.2 and 1.3 (t, 3H), 1.4 (m, 1H), 1.5 (m, 1H), 1.7 (m, 2H), 1.9 (m, 2H), 2.0 (m, 2H), 3.0 (s, 3H), 3.3 (m, 4H), 3.5 (d, 2H), 3.8 (d, 2H), 3.9 and 4.8 (m, 1H), 7.3 (m, 5H), 7.5 (m, 2H), 7.9 (m, 2H); MS: 441 (MH⁺).


Step 2: Preparation of title compound

To a solution of *N*-(8-Benzyl-8-aza-bicyclo[3.2.1]oct-3-yl-exo)-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide (0.37g, 0.85mmol) in ethanol (20mL) was added 20% palladium hydroxide on carbon (0.04g) and the resulting mixture was stirred under an atmosphere of hydrogen for 2 days. The catalyst was removed by filtration and the resulting

solution was adsorbed onto silica. The residue was purified by silica gel chromatography (eluent: DCM to 10% methanol and 1% 0.88 ammonia in DCM) to afford the sub-titled compound as an oil (0.1g); NMR (CDCl₃): 1.1 and 1.2 (t, 3H), 1.3 (m, 1H), 1.4 (m, 2H), 1.7 (m, 5H), 2.1 (br s, 1H), 3.0 (s, 3H), 3.3 (m, 2H), 3.6 (m, 2H), 3.7 and 3.8 (s, 2H), 3.8 and 4.8 (m, 1H), 7.4 (m, 2H), 7.9 (m, 2H); MS: 351 (MH⁺).

Method E


(S)-3-Phenyl-3-(4,4-difluorocyclohexylcarbonylamino)propanal



Step 1: Preparation of (S)-3-amino-3-phenyl-propionic acid methyl ester hydrochloride

To a solution of (S)-3-Bocamino-3-phenyl-propionic acid (5g, 18.8mmol) in methanol (50mL) was added thionyl chloride (1.5mL, 20.7mmol) dropwise. The resulting mixture was stirred at reflux for 4h then allowed to cool and concentrated. The residue was used directly in the next reaction.

Step 2: Preparation (S)-3-phenyl-3-(4,4-difluorocyclohexylcarbonylamino)propanol



To a solution of (S)-3-amino-3-phenyl-propionic acid methyl ester hydrochloride (3.31g, 15.3 mmol) in DCM (50mL) was added triethylamine (1.71g, 17mmol) and the resulting mixture stirred at 0°C for 10min. Then 4,4-difluorocyclohexane carboxylic acid (2.8g, 17mmol) and diisopropylcarbodiimide (2.5g, 17mmol) were added portionwise and the resulting mixture stirred at room temperature for 16h. The mixture was then washed with water, dried over MgSO₄ and concentrated. Silica gel chromatography (eluent: isohexane to diethyl ether) gave the sub-titled compound as a solid (3.7g). This material was then dissolved in THF under an atmosphere of argon and lithium aluminium hydride (11mL, 1M in THF) was added dropwise at 0°C. After stirring for 15min, the reaction was quenched with

2M NaOH and separated. The organic layer was dried over MgSO₄ purified by silica gel chromatography (eluent: isohexane to ethyl acetate) to afford the sub-titled compound as a solid (1.32g); NMR (CDCl₃): 1.8 (m, 8H), 2.2 (m, 3H), 3.6 (m, 1H), 3.7 (m, 1H), 5.2 (m, 1H), 7.3 (m, 5H); MS: 297 (M⁺).

5

Step 3: Preparation of title compound


To a solution of (*S*)-3-phenyl-3-(4,4-difluorocyclohexylcarbonylamino)propanol (0.17g, 0.56mmol) in DCM (5mL) was added Dess Martin periodinane (0.26g, 0.62mmol) and the resulting mixture was stirred for 1h. The mixture was then washed with 2M NaOH, dried over MgSO₄ and concentrated. The resulting residue was then used directly in the preparation of Example 3.

10


Method F

(*S*)-*N*-{1-[3-Amino-3-(3-fluorophenyl)propyl]piperidin-4-yl}-*N*-ethyl-2-(4-methanesulfonyl-phenyl)acetamide

15



Step 1: Preparation of *trans*-3-fluorocinnamic acid *tert*-butyl ester




To a stirred solution of *trans*-3-fluorocinnamic acid (4.34g, 26.1mmol) in toluene (40mL) at 110°C was added *N,N*-dimethylformamide di-*tert*-butyl acetal (25mL, 104mmol) dropwise over 30 min. The resulting mixture was stirred at reflux for a further 4h. The mixture was then cooled to room temperature and washed with water (50mL), saturated aqueous sodium hydrogen carbonate solution (2 x 100mL) and brine (100mL), dried (MgSO₄) and evaporated. The crude product was purified by Bond Elut (isohexane then 2% ethyl acetate in isohexane) to give the title compound as a liquid (3.7g, 64%).

20


25

Step 2: Preparation of (*S*)-3-[(*R*)-benzyl-(1-phenyl-ethyl)-amino]-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester



To a stirred solution of (*R*)-(+)-*N*-benzyl- α -methylbenzylamine (4.0mL, 19mmol) in THF (20mL) at -78°C was added *n*-butyl lithium (1.6M in hexanes, 12.5mL, 20mmol) and the resulting mixture was allowed to warm to room temperature over 10 min. before re-cooling to -78°C . A solution of *trans*-3-fluorocinnamic acid *tert*-butyl ester (3.74g, 16.8mmol) in THF (20mL) was added and the resulting mixture was stirred at -78°C for 2h then quenched by the addition of saturated aqueous ammonium chloride solution (25mL). After warming to room temperature the organic phase was washed with water (2 x 50mL) and brine, dried (MgSO_4) and evaporated. The crude product was purified by Bond Elut (isohexane then 2% ethyl acetate in isohexane) to give the title compound as a gum (5.85g, 80%); NMR (400MHz, CDCl_3): 1.23 (s, 9H), 1.27 (d, 3H), 2.48 (m, 2H), 3.67 (s, 2H), 3.97 (q, 1H), 4.40 (dd, 1H), 6.93 (ddd, 1H), 7.1-7.4 (m, 13H).


Step 3: Preparation of 3-*tert*-butoxycarbonylamino-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester



A stirred mixture of (*S*)-3-[(*R*)-benzyl-(1-phenyl-ethyl)-amino]-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester (5.39g, 12.4mmol), di-*tert*-butyl dicarbonate (2.98g, 13.7mmol) and 20% palladium hydroxide on carbon (0.59g) in ethanol (100mL) was hydrogenated at 5 Bar at room temperature for 24h. The catalyst was removed by filtration through a pad of Celite® washing through with ethanol. The filtrate was evaporated to give an oil which was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate solution. The organic phase was dried (MgSO_4) and evaporated. The crude product was purified by


Bond Elut (eluting with isohexane then 5% ethyl acetate in isohexane) to give the title compound as an oil (3.63g, 86%); NMR: 1.33 (s, 18H), 2.63 (m, 2H), 4.90 (m, 1H), 7.06 (ddd, 1H), 7.24 (m, 2H), 7.37 (dd, 1H), 7.50 (br d, 1H).

- 5 Step 4: Preparation of (*S*)-[1-(3-fluoro-phenyl)-3-hydroxy-propyl]-carbamic acid *tert*-butyl ester



- To a stirred, ice-cooled solution of 3-*tert*-butoxycarbonylamino-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester (2.46g, 7.25mmol) in THF (35mL) was added lithium
10 aluminium hydride (1M in THF, 7.50mL, 7.50mmol) dropwise over 20min. The resulting mixture was stirred with warming to room temperature for 2h. The reaction was quenched with water (0.275mL) then 15% aqueous sodium hydroxide (0.275mL) and more water (0.825mL) were added with stirring. The resultant precipitate was removed by filtration washing with THF, and the filtrate was dried (MgSO₄) and evaporated. The crude product was
15 purified by Bond Elut (gradient elution, isohexane to 30% ethyl acetate in isohexane) to give the title compound as an oil (1.26g, 65%); NMR: 1.4 (s, 9H), 1.75 (m, 1H), 1.85 (m, 1H), 3.3 (m, 1H), 3.4 (m, 1H), 4.5 (dd, 1H), 4.65 (br m, 1H), 7.1 (m + br s, 3H), 7.35 (m, 2H).


- Step 5: Preparation of (*S*)-[1-(3-fluoro-phenyl)-3-oxo-propyl]-carbamic acid *tert*-butyl ester



20

- To a solution of (*S*)-[1-(3-fluoro-phenyl)-3-hydroxy-propyl]-carbamic acid *tert*-butyl ester (0.85g, 3.2mmol) in DCM (70mL) under argon was added Dess-Martin periodinane (1.48g, 3.5mmol) and the resulting mixture was stirred at room temperature for 2h before the addition of 2M aqueous sodium hydroxide (50mL). The organic layer was dried (MgSO₄) and
25 evaporated to give the title compound (quantitative); NMR: 1.4 (s, 9H), 2.8 (m, 2H), 5.1 (m, 1H), 7.05 (ddd, 1H), 7.15 (m, 2H), 7.35 (m, 1H), 7.5 (br d, 1H), 9.6 (s, 1H).

Step 6: Preparation of (*S*)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-(3-fluoro-phenyl)-propyl]-carbamic acid *tert*-butyl ester




5 To a solution of (*S*)-[1-(3-fluoro-phenyl)-3-oxo-propyl]-carbamic acid *tert*-butyl ester (0.85g, 3.12mmol) in DCM (70mL) and *N*-ethyl-2-(4-methanesulfonyl-phenyl)-*N*-piperidin-4-yl-acetamide (Method A, 1.19g, 3.67mmol) was added glacial acetic acid (one drop) and the resulting mixture was stirred at room temperature for 1h. Sodium triacetoxyborohydride (1.4g, 6.4mmol) was added and the resulting mixture was stirred at room temperature for 18 h. The
10 reaction mixture was quenched with water and the organic phase was washed with sodium hydrogen carbonated solution (saturated aqueous) and water, dried (MgSO₄) and concentrated. The crude product was purified by Bond Elut (ethyl acetate then 8% methanol in ethyl acetate) to give the title compound as a solid (1.00g, 55%); NMR: 1.0 and 1.1 (t, 3H), 1.35 (s, 9H), 1.5 (m, 2H), 1.7 (m, 4H), 1.9 (m, 2H), 2.2 (t, 2H), 2.8 (m, 2H), 3.2 (s, 3H), 3.2 and 3.3 (q, 2H),
15 3.6 and 4.1 (m, 1H), 3.8 and 3.85 (s, 2H), 4.5 (m, 1H), 7.05 (m, 1H), 7.1 (m, 2H), 7.35 (dd, 1H), 7.5 (br d, 1H), 7.5 (d, 2H), 7.85 (d, 2H); LCMS: 576 (MH⁺).

Step 7: Preparation of title compound

To a solution of (*S*)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-(3-fluoro-phenyl)-propyl]-carbamic acid *tert*-butyl ester (1.00g, 1.74mmol)
20 in THF (30mL) and water (0.1mL) was added trifluoroacetic acid (5.0mL) and the resulting mixture was stirred at room temperature for 18h. The mixture was evaporated and the residue dissolved in DCM. This solution was washed with 2M aqueous sodium hydroxide, dried (MgSO₄) and evaporated to give the title compound (0.84g, quantitative); NMR: 1.05 and
25 1.09 (t, 3H), 1.45 and 1.50 (m, 2H), 1.75 (m, 4H), 1.95 (m, 2H), 2.25 (m, 2H), 2.88 (m, 2H), 3.20 (s, 3H), 3.25 and 3.30 (q, 2H), 3.67 and 4.08 (m, 1H), 3.82 and 3.89 (s, 2H), 7.00 (m, 1H), 7.15-7.40 (m, 3H), 7.50 (d, 2H), 7.85 (d, 2H), 8.70 (dd, 1H); MS: 476 (MH⁺).

Method G


(*S*)-*N*-{1-[3-Amino-3-(3,5-di-fluorophenyl)propyl]piperidin-4-yl}-*N*-ethyl-2-(4-methanesulfonyl-phenyl)acetamide dihydrochloride




This was prepared from *trans*-3,5-di-fluorocinnamic acid using a similar sequence of reactions to that used to prepare (*S*)-*N*-{1-[3-amino-3-(3-fluorophenyl)propyl]piperidin-4-yl}-*N*-ethyl-2-(4-methanesulfonyl-phenyl)acetamide from *trans*-3-fluorocinnamic acid (Method F) except that the final partitioning between DCM and 2M aqueous sodium hydroxide was omitted.

Method H

N-[1-((4*S*)-4-Phenyl-4-aminobut-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide dihydrochloride




Step 1: Preparation of (*S*)-4-phenyl-4-Boc-aminobutan-2-one



To a solution of (*S*)-*N*-methyl-*N*-methoxy-3-phenyl-3-Boc-aminopropionamide (Step 1 of Method B, 2.02g, 6.56mmol) in THF (70mL) at -78°C was added methylmagnesium chloride (3M in THF, 21.1mmol) dropwise. The resulting mixture was stirred at -78°C for 30min. before warming to room temperature over 3h. The reaction mixture was added to a vigorously stirred mixture of diethyl ether, ice and 1M aqueous potassium dihydrogen phosphate. The aqueous phase was extracted twice with diethyl ether and the combined organic phases washed with sodium hydrogen carbonate solution (sat. aq.) and brine, dried

(Na₂SO₄) and concentrated giving the title compound as a white solid (1.27g, 74%); NMR (CDCl₃): 1.41 (s, 9H), 2.09 (s, 3H), 2.91 (dd, 1H), 3.03 (m, 1H), 5.08 (m, 1H), 5.37 (br s, 1H), 7.28 (m, 5H); MS: 264.

5 Step 2: Preparation of *N*-[1-((4*S*)-4-phenyl-4-Bocaminobut-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide



To a solution of (*S*)-4-phenyl-4-Boc-aminobutan-2-one (1.25g, 4.75mmol) and *N*-(4-piperidinyl)-*N*-ethyl-4-methanesulfonylphenylacetamide (1.54g, 4.75mmol) in THF/1,2-dichloroethane (1:1, 45mL) was added titanium tetrakisopropoxide (3.1mL, 10.45mmol) at room temperature. The resulting mixture was stirred for 15 min. before the addition of sodium triacetoxyborohydride (1.51g, 7.11mmol). The resulting mixture was stirred for 18h before addition of 2M aqueous sodium hydroxide (30mL). The mixture was diluted with DCM, filtered through Celite®, washed with brine, dried (Na₂SO₄) and concentrated. The residue was purified by BondElut chromatography eluting with a mixture of 1% methanol and 0.05% ammonia in ethyl acetate giving the title compound as a white solid (1.04g); MS: 572.

Step 3: Preparation of title compound

To *N*-[1-((4*S*)-4-phenyl-4-Bocaminobut-2-yl)-4-piperidinyl]-*N*-ethyl-4-methanesulfonylphenylacetamide (194mg, 0.339mmol) was added 5M HCl in methanol (5mL) and the resulting mixture stirred at room temperature for 3h. The mixture was evaporated and the residue azeotroped with toluene and triturated with diethyl ether to give the title compound as a white solid (178mg, 98%); MS: 472.

Many intermediates are known in the art, for example 3,3-di-fluoro-cyclobutane carboxylic acid {William R. Dolbier and Dheya M. Al-Fekri; J. Org. Chem. 52, 1872-1874 (1987)}; (*S*)-3-fluoro-pyrrolidine and (*R*)-3-fluoro-pyrrolidine {Giuseppe Giardina, Giulio Dondio and Mario Grugni; *SYNLETT* (1995), 55-57}; and, 4,4-di-fluoro-cyclohexane

carboxylic acid {Mackenzie A R; Marchington A P; Middleton D S; Meadows S D; WO 97/27185-A1}.

EXAMPLE 4

5 The ability of compounds to inhibit the binding of RANTES or MIP-1 α was assessed by an *in vitro* radioligand binding assay. Membranes were prepared from Chinese hamster ovary cells which expressed the recombinant human CCR5 receptor. These membranes were incubated with 0.1nM iodinated RANTES or MIP-1 α , scintillation proximity beads and various concentrations of the compounds of the invention in 96-well plates. The amount of
10 iodinated RANTES or MIP-1 α bound to the receptor was determined by scintillation counting. Competition curves were obtained for compounds and the concentration of compound which displaced 50% of bound iodinated RANTES or MIP-1 α was calculated (IC₅₀). Certain compounds of formula (I) had an IC₅₀ of less than 50 μ M.


15 Results from this test for certain compounds of the invention are presented in Table IV. In Table IV the results are presented as Pic50 values. A Pic50 value is the negative log (to base 10) of the IC₅₀ result, so an IC₅₀ of 1 μ M (that is 1 x 10⁻⁶M) gives a Pic50 of 6. If a compound was tested more than once then the data below is an average of the probative tests results.

TABLE IV

Table Number	Compound number	Pic50
1	1	8.95
1	2	7.68
1	3	7.76
1	6	8.65
2	1	8.48
3	8	9.15

CLAIMS

1. A compound of formula (I):



wherein:

A is CH₂CH₂ or A is absent;

R¹ is C₃₋₇ cycloalkyl (substituted by one or two fluorine atoms and optionally further substituted by C₁₋₄ alkyl) or N-linked heterocyclyl (substituted by one or two fluorine atoms and optionally further substituted by C₁₋₄ alkyl);

R² is C₃₋₆ alkyl or C₃₋₆ cycloalkyl, or phenyl or heteroaryl either of which is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_n(C₁₋₄ alkyl), nitro, cyano or CF₃;

R^{2a}, R⁴ and R^{4a} are, independently, hydrogen or C₁₋₄ alkyl;

R³ and R^{3a} are, independently, hydrogen or C₁₋₄ alkyl or C₁₋₄ alkoxy;

R⁵ is hydrogen, C₁₋₄ alkyl (optionally substituted by halogen, hydroxy, C₁₋₄ alkoxy, C₃₋₇ cycloalkyl, SH, C₁₋₄ alkylthio, cyano or S(O)_q(C₁₋₄ alkyl)), C₃₋₄ alkenyl, C₃₋₄ alkynyl or C₃₋₇ cycloalkyl;

R⁶ is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C₁₋₂)alkyl, heteroaryl(C₁₋₂)alkyl, phenyl(C₁₋₂ alkyl)NH or heteroaryl(C₁₋₂ alkyl)NH;

wherein the phenyl and heteroaryl rings of any of the foregoing are, unless stated otherwise, independently optionally substituted by halo, cyano, nitro, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_mC₁₋₄ alkyl, S(O)₂NR⁷R⁸, NHS(O)₂(C₁₋₄ alkyl), NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHC(O)NH₂, C(O)NH₂, C(O)NH(C₁₋₄ alkyl), NHC(O)(C₁₋₄ alkyl), CO₂H, CO₂(C₁₋₄ alkyl), C(O)(C₁₋₄ alkyl), CF₃, CHF₂, CH₂F, CH₂CF₃ or OCF₃;


R⁷ and R⁸ are, independently, hydrogen or C₁₋₄ alkyl, or together with a nitrogen or

oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C₁₋₄ alkyl, C(O)H or C(O)(C₁₋₄ alkyl);

m, n and q are, independently, 0, 1 or 2;

or a pharmaceutically acceptable salt thereof or a solvate thereof.

2. A compound as claimed in claim 1 wherein R^{2a} , R^3 , R^{3a} and R^4 are all hydrogen.
3. A compound as claimed in claim 1 or 2 wherein R^{4a} is hydrogen or methyl.
4. A compound as claimed in claim 1, 2 or 3 wherein R^1 is C_{3-7} cycloalkyl (substituted by 1 or 2 fluorine atoms and optionally further substituted by C_{1-4} alkyl).
5. A compound as claimed in claim 1, 2, 3 or 4 wherein R^1 is 4,4-di-fluoro-cyclohexyl, 3,3-di-fluoro-cyclopentyl or 3,3-di-fluoro-cyclobutyl.
6. A compound as claimed in claim 1, 2, 3, 4 or 5 wherein R^2 is phenyl or 6-membered heteroaryl optionally substituted by halogen or CF_3 .
7. A compound as claimed in claim 1, 2, 3, 4, 5 or 6 wherein R^5 is ethyl.
8. A compound as claimed in claim 1, 2, 3, 4, 5, 6 or 7 wherein R^6 is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C_{1-2})alkyl, heteroaryl(C_{1-2})alkyl, phenyl(C_{1-2} alkyl)NH or heteroaryl(C_{1-2} alkyl)NH (for example phenyl or phenylCH₂); wherein the phenyl and heteroaryl rings of R^6 are substituted by $S(O)_2C_{1-4}$ alkyl, and optionally further substituted by one or more of halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_mC_{1-4}$ alkyl, $S(O)_2NR^7R^8$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl)₂, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl), $NHC(O)(C_{1-4}$ alkyl), CO_2H , $CO_2(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ; wherein m , R^7 and R^8 are as defined in claim 1.
9. A process for the preparation of a compound of formula (I) as claimed in claim 1, wherein A is absent, comprising treating a compound of formula (II):



with:

an acid chloride of formula $R^1C(O)Cl$, in the presence of a base and in a suitable solvent; or,

an acid of formula R^1CO_2H , in the presence of a suitable coupling agent, a suitable base and in a suitable solvent.

5

10. A pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, and a pharmaceutically acceptable adjuvant, diluent or carrier.

10

11. A compound of the formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, for use in therapy.

12. A compound of formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, in the manufacture of a medicament for use in therapy.

15

13. A method of treating a chemokine mediated disease state in a warm blooded animal suffering from, or at risk of, said disease, which comprises administering to an animal in need of such treatment a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1.

20

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/SE 03/00480

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 211/58, C07D 451/04, C07D 401/12, C07D 403/12, A61K 31/4468,
A61K 31/46, A61P 29/00, A61P 31/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM. ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02070479 A1 (ASTRAZENECA AB), 12 Sept 2002 (12.09.02), page 9 - page 10, Table I, nos. 4-7, 28, the claims --	1-13
P,X	WO 02076948 A1 (ASTRAZENECA AB), 3 October 2002 (03.10.02) --	1-13
X	EP 1013276 A1 (PFIZER INC.), 28 June 2000 (28.06.00), the claims, the examples --	1-13
X	WO 0187839 A1 (ASTRAZENECA AB), 22 November 2001 (22.11.01), page 32 - page 35, nos. 1, 54, 55, the claims --	1-13

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 May 2003

Date of mailing of the international search report

10-06-2003

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Gerd Strandell/BS
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/SE 03/00480

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 0114333 A1 (ASTRAZENECA UK LIMITED), 1 March 2001 (01.03.01) --	1-13
X	WO 0076513 A1 (MERCK & CO., INC.), 21 December 2000 (21.12.00) --	1-13
X	WO 0190106 A2 (PFIZER LIMITED), 29 November 2001 (29.11.01) -- -----	1-13

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **13**
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Claim 13 relates to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practised on the human or animal body (Rule 39.1(iv)). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds or compositions. These alleged effects must be well defined diseases or conditions. The expression "a chemokine mediated disease state" may relate to a number of different disorders and conditions, which can not be clearly defined by this expression. Thus, the search has mainly been restricted to the diseases mentioned in the description on pages 14 to 17.

INTERNATIONAL SEARCH REPORT

Information on patent family members

29/04/03

International Application No.

PCT/SE 03/00480

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	02070479	A1	12/09/02	AU	2382102 A	01/07/02
				GB	0105077 D	00/00/00
				GB	0115579 D	00/00/00
				SE	0103797 D	00/00/00

WO	02076948	A1	03/10/02	GB	0107228 D	00/00/00

EP	1013276	A1	28/06/00	AP	200102186 D	00/00/00
				AP	200102187 D	00/00/00
				AU	1290400 A	31/07/00
				AU	1675100 A	31/07/00
				BG	105709 A	28/02/02
				BG	105721 A	28/02/02
				BR	9905977 A	14/01/03
				BR	9916585 A	16/10/01
				BR	9917007 A	30/10/01
				CA	2350073 A	06/07/00
				CA	2350573 A	06/07/00
				CN	1331591 T	16/01/02
				CN	1331691 T	16/01/02
				EE	200100344 A	15/10/02
				EE	200100345 A	16/12/02
				EP	1140085 A	10/10/01
				EP	1140920 A	10/10/01
				GB	9828420 D	00/00/00
				HR	20010478 A	30/06/02
				HU	0104795 A	29/04/02
				HU	0104910 A	28/10/02
				IL	143510 D	00/00/00
				IL	143512 D	00/00/00
				JP	2000212159 A	02/08/00
				JP	2002533393 T	08/10/02
				JP	2002533461 T	08/10/02
				NO	20013149 A	23/08/01
				NO	20013183 A	08/08/01
				PL	349091 A	01/07/02
				PL	349495 A	29/07/02
				TR	200101793 T	00/00/00
				TR	200101867 T	00/00/00
				TR	200200938 T	00/00/00
				WO	0038680 A	06/07/00
				WO	0039125 A	06/07/00
				GB	9922702 D	00/00/00

WO	0187839	A1	22/11/01	AU	5898101 A	26/11/01
				BR	0110767 A	11/02/03
				EP	1289957 A	12/03/03
				GB	0011838 D	00/00/00
				NO	20025430 A	18/12/02

WO	0114333	A1	01/03/01	AU	6461600 A	19/03/01
				EP	1212299 A	12/06/02
				JP	2003507456 T	25/02/03
				SE	9902987 D	00/00/00

INTERNATIONAL SEARCH REPORT

Information on patent family members

29/04/03

International Application No.

PCT/SE 03/00480

Patent document cited in search report			Publication date	Patent family member(s)			Publication date
WO	0076513	A1	21/12/00	AU	5473800	A	02/01/01
				US	6506777	B	14/01/03

WO	0190106	A2	29/11/01	AU	5248201	A	03/12/01
				EP	1284974	A	26/02/03
				GB	0014046	D	00/00/00
				NO	20025227	A	31/10/02
				US	2002013337	A	31/01/02
				GB	0015835	D	00/00/00
